



Immune response in rainbow trout against infection with *Flavobacterium psychrophilum*

Henriksen, Maya Maria Mihályi; Kania, Per; Madsen, Lone; Buchmann, Kurt; Dalsgaard, Inger

Publication date:
2011

Document Version
Early version, also known as pre-print

[Link back to DTU Orbit](#)

Citation (APA):
Henriksen, M. M. M., Kania, P., Madsen, L., Buchmann, K., & Dalsgaard, I. (2011). *Immune response in rainbow trout against infection with Flavobacterium psychrophilum*. Abstract from 15th International Conference on Diseases of Fish and Shellfish, Split, Croatia.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Immune response in rainbow trout against infection with *Flavobacterium psychrophilum*

Maya M. M. Henriksen^{*1}, Per Kania², Lone Madsen¹, Kurt Buchmann² and Inger Dalsgaard¹

¹National Veterinary Institute, Technical University of Denmark, Copenhagen, Denmark

²Department of Veterinary Disease Biology, University of Copenhagen, Copenhagen, Denmark

The fish pathogen *Flavobacterium psychrophilum* is one of the main causes of mortality in farmed rainbow trout and other salmonid fish. An infected farm can expect mortality rates around 50-60% in fry and 2-10% in juvenile fish within few weeks. Presently no commercial vaccine exists, and fish farmers control the disease with antibiotics. The pathogen has a limited ability to cause disease in an experimental setting without applying a stressor.

A bath-model using 150mg/L H₂O₂ for 60 minutes as a stressor was used on 1.4g rainbow trout fry in four experimental groups: 1) no H₂O₂/no bath infection, 2) H₂O₂/no bath infection, 3) no H₂O₂/ bath infection and 4) H₂O₂/ bath infection. Bath infections were carried out in 10⁷ CFU/ml *F. psychrophilum* bath solution and control groups were bathed in sterile medium. Samples from all internal organs, head and skin were taken before pathogen exposure and 4 hours, 2 days, 5 days, 8 days and 17 days after exposure. The samples will be used for examining the gene expression in rainbow trout related to infection with *F. psychrophilum* and results will be presented. Furthermore, antibody production will be assessed using ELISA on serum samples. The results will be used in order to optimize vaccination or immune-stimulation.